



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

tance, for even if we ultimately adopt broader genera than we do today, we must know more of the interrelations of the groups of species which are included within them, and which for taxonomic purposes must be distinguished by group names of some kind, even though not recognized nomenclaturally.—W. S.

Chapin on New Birds from the Belgian Congo.¹—Mr. Chapin who accompanied Mr. Herbert Lang on an expedition to the Belgian Congo in the interests of the American Museum of Natural History is now engaged in working up the ornithological collections which they secured during their six years' sojourn (1909 to 1915). These comprise some 6200 skins representing about 600 species, and the present paper is the first publication based upon this material. Mr. Chapin's new species are *Chætura melanopygia* (p. 509), *Apaloderma minus* (p. 510) and *Ceriocleptes* (gen. nov.) *xenurus* (p. 512) all from Avakubi, Ituri District, Belgian Congo. Further publications upon this rich collection will be awaited with interest.—W. S.

Riley on New Birds from China and Japan.²—From material received by the U. S. National Museum from China and Japan, Mr. Riley has described three new forms as follows: *Tetrastes bonasia vicinitas* (p. 16), and *Dryocopus martius silvifragus* (p. 162) from Hakodate, Japan, and *Eophona melanura sowerbyi* (p. 163) from Chang Kow Hsien, Hupeh China.—W. S.

Recent Ornithological Papers by Dabbene.³—In the 'Anales of the Buenos Aires Mus., XXVII,' Mr. Dabbene proposes (p. 76) a new genus *Neophlæotomus* for the woodpecker, known as *Phlæotomus schulzi* Cab. and also describes an allied new species *N. shiptoni* (p. 79) from the province of Cordoba, with a colored plate of the male and female.

In two other short papers he reports *Manacus m. gutturosus*, *Harpiprion cayennensis*, *Dendrocygna discolor* and *Sporophila pileata* from Argentina, all of them being new to the fauna.—W. S.

¹ Descriptions of Three New Birds from the Belgian Congo. By James P. Chapin. Bull. Amer. Mus. Nat. Hist., XXXIV, Art. XVI, pp. 509–513. New York, October 20, 1915.

² Descriptions of Three New Birds from China and Japan. By J. H. Riley. Proc. Biol. Soc. Washington, XXVIII, pp. 161–164, September 21, 1915.

³ Description d'un Nouveau Genre et d'une Nouvelle Espèce de Pic Provenant du Nord-Ouest de la République Argentine. por Roberto Dabbene. An. Mus. Nac. Hist. Nat. Buenos Aires, XXVII, pp. 74–81, July 8, 1915.

Una ave nueva para la Argentina. por Roberto Dabbene. Bol. Soc. Physis. I, No. 7, pp. 435–436, Dec. 1914.

Otras especies de aves nuevas. por Roberto Dabbene. *do.* pp. 532–533, June 10, 1915.